How to give the right amount of IBUPROFEN (also known as Motrin, Advil) is different depending on which type of ibuprofen you plan to give.

Dose: Give every 6 hours if needed, for fever or pain. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

Do NOT use with any other medicine containing ibuprofen.

Weight	Age	Infant's Ibuprofen Drops (50 mg / 1.25 mL)	Children's Liquid Ibuprofen (100 mg / 5 mL)	Children's Ibuprofen Chewable Tablets (50 mg)	Junior Strength Ibuprofen Tablets (100 mg)	Adult's Ibuprofen Tablets (200 mg)
0 to 11 pounds (up to 5 kilograms)	0 to 5 months					
12 to 17 pounds (about 6 to 7 kilograms)	6 to 11 months	1.25 mL	2.5 mL			
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	1.875 mL	4 mL			
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	2.5 mL	5 mL → 5mL -12.5 mL -10 mL -7.5 mL -12.5 mL	2 tablets		
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	3.75 mL	-15 mL -12.5 mL -10 mL - 55 mL -7.5 mL	3 tablets		
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	5 mL	10 mL → 15 mL - 12.5 mL 16 mL - 12.5 mL 16 mL - 12.5 mL	4 tablets	2 tablets	1 tablet
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years		-15 mL	5 tablets	2 ½ tablets	1 tablet
72 to 95 pounds (about 33 to 43 kilograms)	11 years		15 mL → (5.00 / 7.5 m)	6 tablets	3 tablets	1 ½ tablets
96 pounds or more (44 kilograms or more)	12 years or older		10 mL → 15 mL-12.5 mL sinL-12.5 mL sinL-12.5 mL sinL-12.5 mL sinL-12.5 mL sinL-12.5 mL	8 tablets	4 tablets	2 tablets
			10 mL + 10 mL = 20 mL		<u> </u>	